

A Cascade of Genes Related to Waardenburg Syndrome

Masayoshi Tachibana

Saitama Cancer Center, Research Institute, Saitama, Japan

On some occasions, mutations of a gene cause different syndromes that may have similar phenotypes. For example, mutations of the *MITF* gene cause Waardenburg syndrome type 2 (Tassabehji *et al*, 1994; Nobukuni *et al*, 1996) as well as Tietz syndrome (Smith *et al*, 1997). On other occasions, mutations of different genes cause an identical syndrome. Molecular analyses of these genes may provide a good opportunity to not only understand such syndromes themselves but also the biologic aspects of cells relevant to these syndromes. By analyzing the genes for Waardenburg syndrome, we showed that *PAX3*,

the gene responsible for Waardenburg syndrome type 1, regulates *MITF*, the gene responsible for Waardenburg syndrome type 2. Such epistatic relationships have been shown between other genes related to Waardenburg syndrome, and likely to construct a cascade. This paper proposes such a cascade, one that involves genes for *PAX3*, *MITF*, human *MyoD*, *MYF5*, *c-MET*, *c-KIT*, tyrosinase, *TRP-1*, human *QNR-71*, *SOX10*, *EDNRB*, and *EDN3*. **Key words:** *EDNRB/MITF/PAX3/SOX10*. *Journal of Investigative Dermatology Symposium Proceedings* 4:126-129, 1999

We have previously described transgenic mice, named *VGA-9*, with an insertional mutation (Tachibana *et al*, 1992). Phenotypes of homozygotes of these mice, i.e., white coat color, microphthalmia, and hearing impairment, are reminiscent of Waardenburg syndrome (WS), whose major symptoms are leucodermia, white forelock, heterochromia irides, and hearing impairment. Skin, hair follicle, eye, and cochlea of mammals contain melanocytes. Thus, the major symptoms of WS individuals and *VGA-9* mice may be due to lack of melanocytes in these tissues. We examined *VGA-9* with special interest in their melanocyte pathology. Indeed, morphologic analysis revealed that homozygous *VGA-9* are lacking in melanocytes in affected tissues, i.e., skin, eye, and cochlea. Cochleae of mammals contain melanocytes as intermediate cells of the stria vascularis, which play an important role for production/maintenance of endolymph and thus for hearing function. Lack of stria intermediate cells caused degeneration of not only stria vascularis but also, successively, the organ of Corti that contains mechanoreceptor hair cells. Thus, all of the revealed phenotypes of *VGA-9* and WS2 can be explained by a lack of melanocytes.

These observations are consistent with the notion that the genes responsible for *VGA-9* and WS2 are involved in melanocyte differentiation/survival. Intriguingly, the murine *microphthalmia* (*mi*) gene, which is mutated in *VGA-9* and in a series of classical "microphthalmia" mice (Hodgkinson *et al*, 1993; Steingrímsson *et al*, 1994), and its human homolog *MITF* (*microphthalmia-associated transcription factor*) gene, which is assigned to chromosome 3p14.1-p12.3 (Tachibana *et al*, 1994) and is mutated in WS2 (Tassabehji *et al*, 1994; Nobukuni *et al*, 1996), encode a protein with basic-helix-loop-helix-leucine-zipper structure. Proteins with such a structure are often transcription

factors that are involved in cell differentiation. To test the hypothesis that *MITF* is involved in melanocyte differentiation, we transfected NIH-3T3 fibroblasts with a *MITF* cDNA inserted in an expression vector (Tachibana *et al*, 1996; Tachibana, 1997), and the transfection converted NIH-3T3 cells into cells with melanocytic characteristics. These cells were dendritic as melanocytes and expressed tyrosinase, tyrosinase-related protein 1 (*TRP-1*), and *TRP-2*, enzymes that catalyze melanin synthesis, although these cells did not synthesize melanin probably due to mutations of the tyrosinase gene in NIH-3T3 cells.

Besides *MITF*, four genes have been shown to be responsible for WS. *PAX3*, the first identified gene for WS, is a human homolog of mouse *Pax-3*, whose mutations were found in *splotch* mice. Mutations of *PAX3* were originally found in WS1 families, whose symptoms include distopia canthorum (lateral displacement of inner canthi of eyes) in addition to the previously mentioned major WS symptoms. Next, *PAX3* mutations were found in families of WS3, whose symptoms include deformity of upper limbs besides the major symptoms. Because distopia canthorum and upper limb deformity cannot be explained by melanocytes anomaly, and because *PAX3* encodes a transcription factor with paired domain and homeodomain, it is likely that *PAX3* is involved in differentiation of cells other than melanocytes. On the other hand, it has not been explained how *PAX3* mutations cause symptoms resulting from melanocyte anomaly in WS1. Two possibilities were considered: the first is that *PAX3* directly differentiates melanocytes; the second is that *PAX3* differentiates melanocytes through transactivation of *MITF*. Expression of *PAX3/Pax-3* in melanocytes is consistent with both of these possibilities; however, we preferred the second possibility because *Pax-3* is expressed earlier and more widely than *mi* in neural crest. This paper describes how we proved our hypothesis that *PAX3* transactivates the *MITF* gene, and failure of the transactivation due to *PAX-3* mutations causes WS1 symptoms related to melanocytes anomaly. Other epistatic relationships of WS genes and protein-protein interaction between their gene products is also discussed.

MATERIALS AND METHODS

Cloning of the *MITF* promoter The approximately 2.3 kb promoter region of *MITF* was isolated from a P1 clone containing the first exon of the *MITF*

Manuscript received April 27, 1999; accepted for publication June 15, 1999
Reprint requests to: Dr. Masayoshi Tachibana, Saitama Cancer Center Research Institute, 818 Komuro, Ina, Saitama 362-0806, Japan.
E-mail: mtachiba@cancer-c.pref.saitama.jp

Abbreviations: EMSA, electrophoretic mobility shift assay; *mi*, *microphthalmia*; *MITF*, *microphthalmia-associated transcription factor*; *TRP-1*, tyrosinase-related protein 1; WS, Waardenburg syndrome.




Figure 1. PAX3 transactivates the *MITF* promoter. The numbers shown at each reporter constructs denote the boundaries upstream and downstream of the initiation site (+1). (a) The *MITF* promoter is efficiently activated in melanoma cells (624-mel) but not in HeLa cells. (b) Co-transfected PAX3 cDNA transactivates the promoter constructs in HeLa cells. (c) The region between position -263 and -238 of the *MITF* promoter is critical for the promoter activity. Deletion and mutation of this region results in loss of promoter activity. Eight copies of this region fused to the basal *MITF* promoter is also activated. (Reproduced from Watanabe *et al.*, 1998 with permission.)

gene. This region was cloned into the luciferase reporter plasmid pGL2-Basic and termed pMITF-2256. Deletion constructs pMITF-382, pMITF-263, pMITF-238, pMITF-382del(260-245), and pMITF-382mut were generated by restriction enzymes or polymerase chain reaction (PCR)-based method.

Cloning of PAX3 cDNA Human PAX3 cDNA containing a Kozak sequence and an entire coding region was generated by reverse transcriptase-polymerase chain reaction (RT-PCR) using RNA from melanoma cells. Three mutations of PAX3 cDNA were designed from reported WS1 mutations and generated by site-directed mutagenesis. The wild-type or mutant PAX3 cDNA were cloned into an expression vector pCEV27.

Reporter assay Approximately 60% confluent cells seeded 1 d before in six-well plates were cotransfected using LipofectAmine (Gibco BRL) reagents as the reporter plasmid (0.5 μ g), effector plasmid (0.5 μ g), and β -galactosidase expression vector PCH110 (0.2 μ g, Pharmacia). After 24 h, cells were lysed and assayed for luciferase activity.

Electrophoretic mobility shift assay (EMSA) Wild-type and mutated PAX3 proteins were derived from *in vitro* transcription/translation reactions. Equivalent amounts of proteins, as determined by [35 S]-methionine labeling, were incubated with end-labeled oligonucleotide of the *MITF* promoter sequence from nucleotide -265 to -236 (approximately 4×10^4 cpm). After incubation the reaction mixture was loaded on a polyacrylamide gel and then run. Details of these procedures were fully described in the original paper (Watanabe *et al.*, 1998).

RESULTS AND DISCUSSION

Approximately 2.3 kb of the *MITF* 5'-upstream region (nucleotides -2256 to +96; transcription site 1) was cloned from a P1 library (GenBank accession number: BankIt: 153821; AF034755). This region and a shorter region (nucleotides -382 to +96) fused to the luciferase gene were activated strongly when transfected in human melanoma cells but not in HeLa cells, as judged by luciferase activity (Fig 1a). We thought the difference was due to the fact that PAX3 is expressed in melanoma cells but not in HeLa cells. Consistent with this notion, ectopic expression of PAX3 in HeLa cells transactivated the *MITF* promoter (Fig 1b). Furthermore, we found a consensus sequence for binding to the homeodomain (ATTAAT) and paired domain (GGAAC) between position -263 and -238. *In vitro* translated/transcribed PAX3 protein was found to bind the promoter sequence around this consensus sequence specifically as judged by EMSA (data not shown). Promoter constructs that do not contain this sequence were not activated in melanoma cells (Fig 1c).

Next we examined the effect of PAX3 mutations observed in WS1 on *MITF* transactivation using reporter assay and EMSA. We selected

three PAX3 mutations that usually occur in the paired domain or homeodomain (Fig 2); Pro50Leu, Arg271Gly, Try266Cys. All of these PAX3 mutations failed to bind or transactivate the *MITF* promoter (Fig 3). These findings show that *MITF* is a target gene of PAX3 and provide a molecular basis for the hypopigmentation and hearing impairment of individuals with WS1.

Pax-3 deficient *splotch* mice (Bober *et al.*, 1994) and PAX3 deficient WS type 3 (WS3) individuals (Hoth *et al.*, 1993) display limb muscle hypoplasia. *Pax-3* plays an important role in appropriate migration of myoblasts to the limb bud (Bober *et al.*, 1994; Goulding *et al.*, 1994; Williams and Ordahl, 1994). Thus, PAX3/*Pax-3* likely regulates genes involved in muscle differentiation. Product of an oncogene, *c-met*, is also required for limb muscle development (Bladt *et al.*, 1995), and this gene encodes the receptor for hepatocyte growth factor of a receptor tyrosine kinase family (Bottaro *et al.*, 1991). Epstein *et al.* (1996) revealed that *c-met* is a target gene of *Pax-3*. First, they showed reduction of *c-met* expression in *splotch* mice. Next, they showed Pax3 can stimulate *c-met* expression in cultured cells. They found putative Pax3 DNA-binding sites in the human *c-MET* promoter. Muscle differentiation and development are dependent on myogenic genes, *MyoD*, *Myf5*, *MyoD*, and *myogenin*. Recently, it was shown that infection of embryonic tissue with a retrovirus encoding Pax-3 is sufficient to induce expression of *MyoD*, *Myf-5*, and *myogenin* (Maroto *et al.*, 1997). Consistent with this finding, investigation of *splotch/Myf-5*^{-/-} double mutant revealed that MyoD acts genetically downstream of Pax-3 and Myf-5 (Tajbakhsh *et al.*, 1997).

Our previous finding of tyrosinase and TRP-1 induction in permanent *MITF* transfectants (Tachibana *et al.*, 1996; Tachibana, 1997) implies that genes for these enzymes are downstream genes of *MITF*. To support this notion it was shown that promoters of the *tyrosinase* and *TRP-1* genes were transactivated by *MITF* through binding with an M-box in these promoters (for review see Shibahara *et al.*, 1998). In contrast, the promoter of the *TRP-2* gene was not transactivated by *MITF* despite the existence of an M-box in the promoter (Yasumoto *et al.*, 1997). Consistent with this finding, we were able to induce melanocytic characteristics by *MITF* in fibroblasts that constitutively express TRP-2 but not in other fibroblasts that do not express it (Tachibana *et al.*, 1994). Others found that *TRP-2* expression did not depend on *mi* in retinal pigment cell (Opdecamp *et al.*, 1997).

The *c-kit* promoter is transactivated by *mi*, and overexpression of *mi* rescues the low *c-kit* expression in mast cells lacking wild-type *mi* (Tsujimura *et al.*, 1996). *Mi* upregulated *c-Kit* expression in melanocyte precursor cells of neural crest, although onset of *c-kit* was independent




Figure 2. Diagram of the PAX3 gene showing location of mutations detected. Each box indicates an exon. Numbers in parentheses denote the numbers of deleted nucleotides. Non-truncating mutations are expected to encode a full-length PAX3 with a substitution of an amino acid at the paired or homeodomain. S, splice-site mutation (modified from Tassabehji *et al*, 1995; mutations cited by Baldwin *et al*, 1995; Tassabehji *et al*, 1995; Watanabe *et al*, 1998 are shown).




Figure 3. Mutant PAX3 failed to bind and transactivate the MITF promoter. (a) EMSA: PAX3[P50 L] (predicting the substitution of an amino acid in the paired domain) and PAX3[R271G]/PAX3[W266C] (predicting the substitution of an amino acid in the homeodomain) fail to bind the consensus sequence (ATTAATAC-TACTGGAAC) in the MITF promoter sequence, whereas wild-type PAX3 does bind the sequence. Arrowheads identify higher (H) and lower (L) DNA-protein complexes, and the free probe (F). The higher complex (H) is sequence specific and contains PAX3 protein, whereas the lower complex (L) is not sequence specific and does not contain PAX3 (not shown). (b) Reporter assay: Co-transfection of mutated PAX3 cDNA, designed for mutations found in WS1, failed to transactivate the MITF promoter, whereas wild-type PAX3 cDNA did transactivate. (Reproduced from Watanabe *et al*, 1997 with permission.)

of *mi* (Opdecamp *et al*, 1997). Stimulation of c-Kit receptor in melanoma cells activated MAP kinase, which in turn phosphorylated Mi by recruiting p300/CBP and consequently upregulated the tyrosinase gene (Hemesath *et al*, 1998; Price *et al*, 1998). These findings suggest that *c-kit* is a downstream gene of *mi* and interaction of Mi and that c-Kit is crucial for melanocyte development. Mutations of *c-KIT* cause phenotypes similar to WS: piebaldism associated with deafness in human (Spitz and Beighton, 1998). Another gene shown to be downstream of *mi* is the quail *QNR-71* gene, which is expressed specifically in pigment layers and encodes a protein homologous to melanosomal matrix protein. The promoter of this gene is transactivated by two b-HLZip proteins, i.e., Mi and c-MYC (Turque *et al*, 1996).

WS associated with Hirschprung disease, i.e., aganglionic megacolon, is classified as WS type 4 (WS4). Mutations of three genes have been identified in WS4: genes for endothelin-B receptor or endothelin 3 and the *SOX10* gene. Disruption of the murine endothelin-3 (*Edn3*) gene and the endothelin-receptor (*Ednrb*) gene results in aganglionic megacolon and spotted coat color in mice (Baynash *et al*, 1994; Hosoda

et al, 1994). Human mutations of *EDN3* (Edrey *et al* 1996; Hofstra *et al*, 1996) and *EDNRB* (Puffenberger *et al*, 1994) are associated with WS4 families. Mutations of murine *Sox10* or human *SOX10*, encoding a transcription factor with the HMG DNA-binding motif, were found, respectively, in *Dom* mice with megacolon (Pingault *et al*, 1998) and WS4 families (Southard-Smith *et al*, 1998). Thus, interaction of *EDN3*/*Edn3* and *EDNRB*/*Ednrb*, and activation of *SOX10*/*Sox10*, seem to be essential for development of epidermal melanocytes and enteric neurons; however, the question of how these gene products fit in the cascade of WS genes remains to be answered. While *Edn3* enhanced the number of melanocytes in wild-type neural crest cultures (Reid *et al*, 1996), it did not do so in those from *mi*-mutated mice (Opdecamp *et al*, 1997). These findings are consistent with the notion that *Ednrb* is the downstream of *mi*, and, consequently *Ednrb* is not expressed in neural crest cells of *mi*-mutant mice. Because *Ednrb* positive cells are expressed in migrating neural crest cells of 10.5-d.p.c. *Sox10*^{Dom}/*Sox10*^{Dom} embryos (Southard-Smith *et al*, 1998), *Ednrb* seems to be independent of *Sox10*. Furthermore, since *Ednrb* and *TRP-1*-




Figure 4. A cascade of genes and/or their products related to Waardenburg syndrome. Ovals represent transcription factors, and rectangles represent other types of proteins; those responsible for WS are in black. PAX3, human MyoD, MYF5, and c-MET are involved in myocyte differentiation/proliferation, whereas PAX3, MITF, SOX10, c-KIT, tyrosinase, TRP-1, human QNR-71, EDNRB, and EDN3 are involved in melanocyte differentiation/proliferation. It is unlikely that SOX10 regulates EDNRB, and thus the downstream genes of SOX10 remains to be elucidated.

positive cells were virtually absent in neural crest of the 11.5-d.p.c embryos (Southard-Smith *et al*, 1998), *Sox10* seems to determine the neural crest cell fate.

To summarize, three transcription factor genes, *PAX3*, *MITF*, and *SOX10*, have been shown to be responsible for WS. By epistatic relationship, the *PAX3* gene most likely regulates the human *MyoD*, *MYF5*, and *c-MET* genes, whereas the *MITF* gene regulates genes for c-KIT, tyrosinase, TRP-1, and human QNR-71. In addition, we determined the epistatic relationship of *PAX3* and *MITF* (Watanabe *et al*, 1998). Further studies will determine whether there is an epistatic relationship between *SOX10* and *MITF* or *PAX3*. Completion of cascade-like network among genes and/or gene products related to WS (Fig 4) will help us understand the mechanism of the manifestation of WS symptoms.

I thank my colleagues Drs A Watanabe and K Takeda, without whose help the present work would not have been accomplished.

REFERENCES

- Baldwin CT, Hoth CF, Macina RA, Milunski A: Mutations in PAX3 that cause Waardenburg syndrome type I. *Am J Med Genet* 58:115-122, 1995
- Baynash AG, Hosoda K, Giald A, Richardson JA, Emoto N, Hammer RE, Yanagisawa M: Interaction of endothelin-3 with endothelin-B receptor is essential for development of epidermal melanocytes and enteric neurons. *Cell* 79:1277-1285, 1994
- Bladt F, Riethmacher D, Isenmann S, Aguzzi A, Birchmeier C: Essential role for the c-met in the migration of myogenic precursor cells into the limb bud. *Nature* 376:768-771, 1995
- Bober E, Franz T, Arnold H, Gruss P, Tremblay P: Pax-3 is required for the development of limb muscles: a possible role for the migration of dermomyotomal muscle progenitor cells. *Development* 120:603-612, 1994
- Bottaro DP, Ruin J, Faletto DL, Chan AML, Kmiec TE, Vande Woude GF, Aaronson SA: Identification of the hepatocyte growth factor receptor as the c-met proto-oncogene product. *Science* 251:802-804, 1991
- Edery P, Attie T, Amiel J, *et al*: Mutation of the endothelin-3 gene in the Waardenburg-Hirschprung disease (Shah-Waardenburg syndrome). *Nature Genet* 12:445-447, 1996
- Epstein JA, Shapiro DN, Cheno J, Lam PYP, Maas RL: Pax3 modulates expression of c-Met receptor during limb muscle development. *Proc Natl Acad Sci USA* 93:4213-4218, 1996
- Goulding M, Lumsden A, Paquette AJ: Regulation of Pax-3 expression in the dermomyotome and its role in muscle development. *Development* 120:957-971, 1994
- Hemesath TJ, Price ER, Takemoto C, Badalian T, Fisher DE: MAP kinase links the transcription factor Microphthalmia to c-Kit signaling in melanocytes. *Nature* 391:298-301, 1998
- Hodgkinson CA, Moore KJ, Nakayama A, Steingrimsson E, Copeland NG, Jenkins NA, Arnheiter H: Mutations at the mouse microphthalmia locus are associated with defects in a gene encoding novel basic-helix-loop-helix-zipper protein. *Cell* 74:395-404, 1993
- Hofstra RM, Osinga J, Tan-Sindhunata G, *et al*: A homozygous mutation in the endothelin-3 gene associated with a combined Waardenburg type 2 and Hirschprung phenotype (Shah-Waardenburg syndrome). *Nature Genet* 13:395-396, 1996
- Hosoda K, Hammer RE, Richardson JA, Baynash AG, Cheung JC, Giald A, Yanagisawa M: Targeted and natural (piebald-lethal) mutations of endothelin-B receptor gene produce megacolon associated with spotted coat color in mice. *Cell* 79:1267-1276, 1994
- Hoth CF, Milunsky N, Lipsky R, Sheffer SK, Baldwin CT: Mutations in the paired domain of the human PAX3 gene cause Klein-Waardenburg syndrome (WS-III) as well as Waardenburg syndrome I (WS-I). *Am J Hum Genet* 52:455-462, 1993
- Marato M, Reshef RAE, Munsterberg Koester S, Goulding M, Lassar A: Ectopic Pax-3 activates *MyoD* and *Myf-5* expression in embryonic mesoderm and neural tissue. *Cell* 89:139-148, 1997
- Nobukuni Y, Watanabe A, Takeda K, Skarka H, Tachibana M: Analyses of loss-of-function mutations of the MITF gene suggest that haploinsufficiency is a cause of Waardenburg syndrome type 2A. *Am J Hum Genet* 59:76-83, 1996
- Opdecamp K, Nakayama A, Nguyen M-TT, Hodgkinson CA, Pavan WJ, Arnheiter H: Melanocyte development in vivo and in neural crest cultures: crucial dependence on the Mitf basic-helix-loop-helix-zipper transcription factor. *Development* 124:2377-2386, 1997
- Pingault V, Bondurand N, Kuhlbrod K, *et al*: SOX10 mutations in patients with Waardenburg-Hirschprung disease. *Nature Genet* 18:171-173, 1998
- Price ER, Ding HF, Badalian T, *et al*: Lineage-specific signaling in melanocytes. C-kit stimulation recruits p300/CBP to microphthalmia. *J Biol Chem* 273:17983-17986, 1998
- Puffenberger EG, Hosoda K, Washington SS, Nakao K, deWit D, Yanagisawa M, Chakravart A: A missense mutation of the Endothelin-B receptor gene in multigenic Hirschprung's disease. *Cell* 79:1257-1266, 1994
- Reid K, Turnley AM, Maxwell GD, Kurihara Y, Kurihara H, Bartlett PF, Murphy M: Multiple roles for endothelin in melanocyte development: regulation of progenitor number and stimulation of differentiation. *Development* 122:3911-3919, 1996
- Shibahara S, Yasumoto K, Takahashi K: Genetic regulation of the pigment cell. In: Nordlund JJ, Boissy RE, Hearing VJ, King RA, Ortonne J-P (eds). *The Pigment System, Physiology and Pathology*. New York: Oxford University Press, 1998, pp. 251-273
- Southard-Smith EM, Kos L, Pavan WJ: *Sox10* mutation disrupts neural crest development in Dom Hirschprung mouse model. *Nature Genet* 18:60-64, 1998
- Spitz RA, Beighton P: Piebaldism with deafness; molecular evidence for an expanded syndrome. *Am J Med Genet* 75:101-103, 1998
- Steingrimsson E, Moore KJ, Lamoreux ML, *et al*: Molecular basis for mouse *microphthalmia* (*mi*) mutations helps explain their developmental and phenotypic consequences. *Nature Genet* 8:256-263, 1994
- Tachibana M: Evidence to suggests that expression of *MITF* induces melanocyte differentiation and haploinsufficiency of MITF causes Waardenburg syndrome type 2A. *Pigment Cell Res* 10:25-33, 1997
- Tachibana M, Hara Y, Vyas D, Hodgkinson C, Fex J, Grundfast K, Arnheiter H: Cochlear disorder associated with melanocyte anomaly in mice with a transgenic insertional mutation. *Mol Cell Neurosci* 3:433-445, 1992
- Tachibana M, Perez-Jurado LA, Nakayama A, *et al*: Cloning of *MITF*, the human homolog of the mouse microphthalmia gene and assignment to chromosome 3p14.1-p12.3. *Hum Mol Genet* 3:553-557, 1994
- Tachibana M, Takeda K, Nobukuni Y, *et al*: Ectopic expression of *MITF*, a gene for Waardenburg syndrome type 2, converts fibroblasts to cells with melanocyte characteristics. *Nature Genet* 14:50-54, 1996
- Tajbakhsh S, Rocancourt D, Cossu G, Buckingham M: Redefining the genetic hierarchies controlling skeletal myogenesis: Pax-3 and Myf-5 act upstream of MyoD. *Cell* 89:127-138, 1997
- Tassabehji M, Newton VE, Read AP: Waardenburg syndrome type 2 caused by mutations in the human microphthalmia (*MITF*) gene. *Nat Genet* 8:251-255, 1994
- Tassabehji M, Newton VE, Liu X-Z, *et al*: The mutational spectrum in Waardenburg syndrome. *Hum Mol Genet* 11:2131-2137, 1995
- Tsujimura T, Morii E, Nozaki M, *et al*: Involvement of transcription factor encoded by the *mi* locus in the expression of c-kit receptor tyrosine kinase in cultured mast cells of mice. *Blood* 88:1225-1233, 1996
- Turque N, Denhez F, Martin P, Planque N, Bailly M, Bègue Stèhelin D, Saule S: Characterization of a new melanocyte-specific gene (*QNR-71*) expressed in v-myc-transformed quail neuroretina. *EMBO* 15:3338-3350, 1996
- Watanabe A, Takeda K, Ploplis B, Tachibana M: Epistatic relationship between Waardenburg syndrome genes *MITF* and *PAX3*. *Nature Genet* 18:283-286, 1998
- Williams BA, Ordahl CP: Pax-3 expression in segmental mesoderm marks early stages in myogenic cell specification. *Development* 120:785-789, 1994
- Yasumoto K, Yokoyama K, Takahashi K, Tomita Y, Shibahara S: Functional analysis of microphthalmia-associated transcription factor in pigment cell-specific transcription of the human tyrosinase family genes. *J Biol Chem* 272:503-509, 1997